		Residence		Education					
Key Indicators for Uttar Pradesh							10 years		
from NFHS-3	NFHS-3 (2005-06)	Urban	Rural	No education <sup>4</sup>	<5 years complete <sup>4</sup>	5-9 years complete <sup>4</sup>	complete and above <sup>4</sup>	NFHS-2 (1998-99)	NFHS-1 (1992-93)
Marriage and Fertility	(2003-00)	Orban	Kulai	education	complete	complete	and above	(1990-99)	(1992-93)
1. Women age 20-24 married by age 18 (%)	58.6	32.1	67.9	77.8	61.7	56.4	23.0	64.3	na
2. Men age 25-29 married by age 21 (%)	51.4	29.5	60.8	63.7	60.5	56.5	36.8	na	na
3. Total fertility rate (children per woman)	3.8	3.0	4.1	4.6	3.3	3.3	2.4	4.1	na
4. Women age 15-19 who were already mothers or pregnant at the time of the survey (%)	14.3	7.8	16.3	26.4	11.4	12.3	4.3	na	na
5. Median age at first birth for women age 25-49	19.4	20.3	19.1	18.7	19.5	19.8	22.6	18.8	na
6. Married women with 2 living children wanting no more children (%)	65.5	76.9	60.1	54.1	(38.8)	70.7	83.3	43.7	na
6a. Two sons	73.7	78.3	71.4	66.9	*	76.7	84.6	65.2	na
6b. One son, one daughter	72.2	84.2	66.5	57.4	*	78.7	90.2	43.8	na
6c. Two daughters	30.8	51.9	21.3	23.8	*	32.2	52.2	16.5	na
Family Planning (currently married women, age 15–49)									
Current use 7. Any method (%)	43.6	56.3	39.7	39.7	40.1	46.5	58.1	27.1	na
8. Any modern method (%)	29.3	42.4	25.3	25.8	28.2	30.6	44.2	20.8	na na
8a. Female sterilization (%)	17.3	18.7	16.8	18.5	17.3	16.1	13.5	14.1	na
8b. Male sterilization (%)	0.2	0.5	0.1	0.1	0.4	0.1	0.2	0.5	na
8c. IUD (%)	1.4	3.2	0.8	0.7	0.9	1.2	5.0	0.9	na
8d. Pill (%)	1.7	3.2	1.3	1.0	1.3	2.5	3.8	1.2	na
8e. Condom (%)	8.6	16.6	6.2	5.3	7.5	10.6	21.4	4.0	na
Unmet need for family planning									
9. Total unmet need (%)	21.2	15.1	23.1	22.0	22.8	21.4	16.5	25.4	na
9a. For spacing (%)	9.1	5.8	10.1	8.3	11.0	11.5	8.3	11.8	na
9b. For limiting (%)	12.1	9.3	13.0	13.7	11.8	9.9	8.2	13.6	na
Maternal and Child Health									
Maternity care (for births in the last 3 years)	26.2	40.0	22.6	171	24.0	22.0	(11	146	
10. Mothers who had at least 3 antenatal care visits for their last birth (%) 11. Mothers who consumed IFA for 90 days or more when they were pregnant with their	26.3	40.9	22.6	17.1	24.0	33.0	64.1	14.6	na
last child (%)	8.7	16.4	6.7	3.8	4.7	13.6	26.6	na	na
12. Births assisted by a doctor/nurse/LHV/ANM/other health personnel (%) <sup>2</sup>	29.2	50.5	23.8	19.2	24.6	37.5	68.8	21.8	na
13. Institutional births (%) <sup>2</sup>	22.0	39.9	17.5	13.6	19.6	26.6	59.2	15.2	na
14. Mothers who received postnatal care from a doctor/nurse/LHV/ANM/other health									
personnel within 2 days of delivery for their last birth (%) <sup>2</sup>	14.3	31.1	10.0	7.0	16.0	17.7	46.2	na	na
Child immunization and vitamin A supplementation <sup>2</sup> 15a. Children 12-23 months fully immunized (BCG, measles, and 3 doses each of									
polio/DPT) (%)	23.0	33.0	20.5	13.7	(30.8)	34.0	52.6	20.2	na
15b. Children 12-23 months who have received BCG (%)	61.0	66.6	59.6	50.7	(74.8)	75.8	88.2	56.5	na
15c. Children 12-23 months who have received 3 doses of polio vaccine (%)	87.6	89.3	87.1	85.4	(77.9)	90.7	96.1	41.3	na
15d. Children 12-23 months who have received 3 doses of DPT vaccine (%)	30.0	38.7	27.9	20.2	(30.8)	41.9	62.3	32.7	na
15e. Children 12-23 months who have received measles vaccine (%)	37.7	49.2	34.8	27.0	(49.6)	50.7	70.3	33.5	na
16. Children age 12-35 months who received a vitamin A dose in last 6 months (%)	8.7	10.5	8.2	6.4	17.5	10.7	15.3	na	na
Treatment of childhood diseases (children under 3 years) <sup>2</sup>	12.0	15.0	44.0	0.5	*	12.4	24.7	45.0	
17. Children with diarrhoea in the last 2 weeks who received ORS (%) 18. Children with diarrhoea in the last 2 weeks taken to a health facility (%)	12.0 60.2	15.0 66.3	11.2 58.7	9.5 59.5	*	12.4 61.6	24.7 57.5	15.2 63.3	na
19. Children with acute respiratory infection or fever in the last 2 weeks taken to a health	00.2	00.5	30.7	39.3		01.0	37.3	03.3	na
facility (%)	70.4	85.3	66.7	66.7	(82.6)	72.9	80.8	na	na
Child Feeding Practices and Nutritional Status of Children <sup>2, 3</sup>									
20. Children under 3 years breastfed within one hour of birth (%)	7.2	8.5	6.9	5.2	7.9	8.1	15.7	5.7	na
21. Children age 0-5 months exclusively breastfed (%)	51.3	31.8	55.8	55.0	(63.1)	48.0	35.0	na	na
22. Children age 6-9 months receiving solid or semi-solid food and breastmilk (%)	45.5	49.3	44.3	40.2	*	46.9	64.6	na	na
23. Children under 3 years who are stunted (%)	52.4	46.6	53.6	58.2	53.9	47.2	32.0	60.7	na
24. Children under 3 years who are wasted (%)	19.5	16.5	20.3	21.2	25.6	17.3	14.0	16.9	na
25. Children under 3 years who are underweight (%)  Nutritional Status of Ever-Married Adults (age 15-49)	41.6	31.8	43.7	47.8	43.5	34.5	21.9	48.1	na
26. Women whose Body Mass Index is below normal (%)	34.1	23.3	37.2	39.0	32.3	30.0	17.8	36.5	na
27. Men whose Body Mass Index is below normal (%)	32.7	21.9	36.5	42.8	40.5	35.3	18.9	na	na
28. Women who are overweight or obese (%)	11.1	24.6	7.2	7.1	10.0	14.2	25.4	7.3	na
29. Men who are overweight or obese (%)	9.9	22.0	5.7	2.7	4.3	7.9	19.9	na	na
Anaemia among Children and Adults									
30. Children age 6-35 months who are anaemic (%)	85.0	82.5	85.7	86.7	85.1	83.4	80.1	73.8	na
31. Ever-married women age 15-49 who are anaemic (%)	50.9	50.7	50.8	52.7	54.9	48.4	44.6	49.0	na
32. Pregnant women age 15-49 who are anaemic (%)	51.6	51.3	51.7	55.7	*	44.9	40.4	45.8	na
33. Ever-marrried men age 15-49 who are anaemic (%)	24.7	15.7	27.8	29.5	25.1	25.6	19.5	na	na
Knowledge of HIV/AIDS among Ever-Married Adults (age 15-49)									
34. Women who have heard of AIDS (%)	40.0	67.7	31.3	21.4	40.5	63.2	91.9	19.2	na
35. Men who have heard of AIDS (%) 36. Women who know that consistent condom use can reduce the chances of getting	74.3	88.2	69.1	44.6	60.1	80.1	97.7	na	na
HIV/AIDS (%)	27.2	53.1	19.2	11.5	21.9	43.2	78.2	na	na
37. Men who know that consistent condom use can reduce the chances of getting									
HIV/AIDS (%)	66.3	82.4	60.3	34.8	49.9	70.8	93.4	na	na
Women's Empowerment	25 =		2.4	0 = -	26 =	00.5	25.5		
38. Currently married women who usually participate in household decisions (%) 39. Ever-married women who have ever experienced spousal violence (%)	33.7	41.1	31.4	35.2	29.7	29.9	33.2	na	na
ээ. Ever-married women wno nave ever experienced spousal violence (%)	42.4	36.0	44.3	48.8	45.0	37.2	18.3	na	na

na : Not available

<sup>()</sup> Based on 25-49 unweighted cases

<sup>\*</sup> Based on fewer than 25 unweighted cases

<sup>&</sup>lt;sup>1</sup> Excludes pregnant women.

 $<sup>^{2}</sup>$  Based on the last 2 births in the 3 years before the survey to ever-married women.

<sup>&</sup>lt;sup>3</sup> Based on WHO standard.

<sup>&</sup>lt;sup>4</sup> For children education refers to mother's education. Children with missing information on the mother's education are not included in the education columns.